





PubMed Ni	ucleotide	Protein	Genome	Stru	cture	PMC	Taxonom	у	●MIM Book
Search PubMed	v for							Go	Clear
	Limits	Previev	//Index His	story	CI	ipboard	Details		
			F-3					-	
	Display	Summary	▼ Show:	20	▼ Sc	ort 🔻	Send to	Text	
C-6		Items 1-	20 of 153				Pag	e <u>1</u>	of 8 Nex
Entrez PubMed	☐1: Kumar NS, Schaefer PA, Lark G, Frieri M. Related Articles, Links Late phase response during nasal challenge: effect of astemizole on leukotriene B4 levels. Allergy Asthma Proc. 1996 Mar-Apr;17(2):93-9. PMID: 8934801 [PubMed - indexed for MEDLINE]								
PubMed Services	Sanso KB. Effect seasor Ann Al	ne G, Etzel J. of mometa nal allergic lergy Asthma		on ea	M, Mesa arly and 81(5 Pt	arina-Wicki l late phas 1):431-7.	B, Nolop		ted Articles, Lin
Related Resources	Intranasal mometasone furoate reduces late-phase inflammation after allergen challenge. Ann Allergy Asthma Immunol. 2001 Apr;86(4):433-8. PMID: 11345288 [PubMed - indexed for MEDLINE]								
	14: Alvarado-Valdes CA, Blomgren J, Weiler D, Gleich GJ, Reed CE, Field Related Articles, Links EA, Wisniewski ME, Pobiner BF. The effect of fluticasone propionate aqueous nasal spray on eosinophils and cytokines in nasal secretions of patients with ragweed allergic rhinitis. Clin Ther. 1997 Mar-Apr;19(2):273-81. PMID: 9152566 [PubMed - indexed for MEDLINE]								
	5: Jacobi HH, Poulsen LK, Reimert CM, Skov PS, Ulfgren AK, Jones I, Elfman LB, Malling HJ, Mygind N. IL-8 and the activation of eosinophils and neutrophils following nasal allergen challenge. Int Arch Allergy Immunol. 1998 May;116(1):53-9. PMID: 9623510 [PubMed - indexed for MEDLINE]								
	☐ 6: Freeland HS, Pipkorn U, Schleimer RP, Bascom R, Lichtenstein LM, Naclerio RM, Peters SP. Leukotriene B4 as a mediator of early and late reactions to antigen in humans: the effect of systemic glucocorticoid treatment in vivo. J Allergy Clin Immunol. 1989 Mar;83(3):634-42. PMID: 2466880 [PubMed - indexed for MEDLINE]								







PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Senich Public	or . To for	norastani	sole and 1	enkotitienė	1. J. T. 11. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1		Clear	
(A 3. 19.6: 2 3 1	Linus	Previe	w/lindex 出出	islow Cl	rbisodo	. Details	100	
	*							
	No items	found.						


See Details.

Entrez PubMed

PubMed Services

Related Resources

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer



DATE: Monday, February 03, 2003 Printable Copy Create Case

Set Name	<u>Query</u>	Hit Count	Set Name
side by side			result set
DB = USPT, PG	GPB,JPAB,EPAB,DWPI; PLUR=YES; OP=AND	•	
<u>L4</u>	L3 norastemizole	17	<u>L4</u>
<u>L3</u>	antihistamines leukotriene	618	<u>L3</u>
<u>L2</u>	norastemizole montelukast	15	<u>L2</u>
L1	norastemizole leukotriene	19	<u>L1</u>

END OF SEARCH HISTORY







> NCDI	of Medicine NLM	
PubMed N	cleotide Protein Genome Structure PMC Taxonomy OMIM Boo	ks
Search PubMed	▼ for Go Clear	
	Limits Preview/Index History Clipboard Details	
<u> </u>	Display Abstract Show: 20 Sort Send to Text	
Entrez PubMed	☐1: Allergy Asthma Proc 1996 Mar-Apr;17(2):93-9 Related Articles, Lin Late phase response during nasal challenge: effect of astemizole of	
	leukotriene B4 levels.	. ,
	Kumar NS, Schaefer PA, Lark G, Frieri M.	
PubMed Services	Department of Pathology and Surgery, Nassau County Medical Center, State University of New York at Stonybrook, USA.	7 0 0, 50
Related Resources	Nasal provocation (NP) in allergic rhinitis patients can elicit a late phase inflammatory response in which interleukins (IL), leukotrienes (LT), and neutrop have been implicated. Certain antihistamines have been shown to have anti-inflammatory properties. The objective was to determine whether astemizole 10 mgs/day has any anti-inflammatory characteristics. We clinically evaluated 20 patients with allergic rhinitis and measured nasal IL-8 and LTB4 during NP with increasing doses of grass and ragweed antigen in a double-blind placebo-controlle fashion after a 4-week course of treatment. Clinical symptom scores for sneezing pruritus, and rhinorrhea were evaluated. Nasal fluid was examined by ELISA and RIA for IL-8 and LTB4 levels along with neutrophil assessment before NP and at 6, and 8 hours. Symptom scores for nasal sneezing, pruritus, rhinorrhea, and nasa LTB4 levels at 6 and 8 hours and IL-8 at 3, 6, and 8 hours were generally lower in astemizole-treated patients compared to those on placebo. Nasal IL-8 levels corresponded to LTB4 levels at diluent and at 6 hours in the placebo group (P = 0.01). The percentage of neutrophils correlated with LTB4 levels at baseline, coefficient = 0.76, P = 0.02 and at 6 hours, coefficient = 0.62, P = 0.035 in the placebo group. This study is the first to demonstrate an effect of astemizole durin NP on IL-8 and LTB4 levels with a significant correlation of neutrophil numbers untreated patients during the nasal late phase response.	e at deed , , , , , , , , , , , , , , , , ,
	Publication Types:	
	PMID: 8934801 [PubMed - indexed for MEDLINE]	
	Display Abstract ▼ Show: 20 ▼ Sort ▼ Send to Text ▼	